

Citoplastia de aumento. Se realiza cuando las otras alternativas de tratamiento no han funcionado. Ha demostrado ser efectivo en pacientes neurológicos, incluidos EM²³, y aunque existen riesgos asociados, con los cuidados apropiados parece una técnica segura.

Bibliografía

1. Fundación Esclerosis Múltiple. Disponible en: <http://www.fem.es/la-incidencia-de-la-esclerosis-multiple-en-espana-ha-aumentado-un-50-en-los-ultimos-20-anos.aspx>. Actualizado diciembre 2013. [consultado 2/3/18].
2. Nicholas RS, Friede T, Hollis S, Young CA. Anticholinergics for urinary symptoms in multiple sclerosis. *Cochrane Database Syst Rev*. 2009 Jan 21;(1):CD004193. doi: 10.1002/14651858.CD004193.pub2.
3. Del Popolo G, Panariello G, Del Corso F, De Scisciolo G, Lombardi G. Diagnosis and therapy for neurogenic bladder dysfunctions in multiple sclerosis patients. *Neurol Sci*. 2008 Dec;29 Suppl 4:S352-5. doi: 10.1007/s10072-008-1042-y.
4. Betts C, D'Mellow M, Fowler C. Urinary symptoms and the neurological features of bladder dysfunction in multiple sclerosis. *J Neurol Neurosurg Psychiatry* 1993; 56: 245-50.
5. Chancellor MB, Anderson RU, Boone TB (2006) Pharmacotherapy for neurogenic detrusor overactivity. *Am J Phys Med Rehabil* 85:536–545.
6. Diokno A, Ingber M (2006) Oxybutynin in detrusor overactivity. *Urol Clin N Am* 33:439–445.
7. Di Stasi SM, Giannantoni A, Navarra P et al (2001) Intravesical oxybutynin: mode of action assessed by passive diffusion and electromotive administration with pharmacokinetics of oxybutynin and N-desethyl oxybutynin. *J Urol* 166:2232–2236.
8. Elkelini MS, Abuzgaya A, Hassouna MM. Mechanisms Of Action Of Sacral Neuromodulation. *Int Urogynecol J*. 2010;21: S439–46. Authors discuss different ways for SNM to control various types of voiding dysfunction. 13. McClurg D, Ashe RG, Marshall K,
9. De Ridder D, Vermeulen C, Ketelaer P et al (1999) Pelvic floor rehabilitation in multiple sclerosis. *Acta Neurol Belg* 99:61–64.
10. Ishigooka M, Hashimoto T, Hayami S et al (1996) Electrical pelvic floor stimulation: a possible alternative treatment for reflex urinary incontinence in patients with spinal cord injury. *Spinal Cord* 34:411–415.
11. McClurg D, Ashe RG, Marshall K, Lowe-Strong AS (2006) Comparison of pelvic floor muscle training, electromyography biofeedback, and neuromuscular electrical stimulation for bladder dysfunction in people with multiple sclerosis: a randomized pilot study. *Neurourol Urodyn* 25:337–348.
12. Rickwood AM (2002) Assessment and conservative management of the neuropathic bladder. *Semin Pediatr Surg* 11:108–119.
13. Lúcio A, D'ancona CA, Perissinotto MC, McLean L, Damasceno BP, de Moraes Lopes MH. Pelvic Floor Muscle Training With and Without Electrical Stimulation in the Treatment of Lower Urinary Tract Symptoms in Women With Multiple Sclerosis. *J Wound Ostomy Continence Nurs*. 2016 Jul-Aug;43(4):414-9. doi: 10.1097/WON.0000000000000223.
14. Tubaro A, Puccini F, De Nunzio C, Digesu GA, Elneil S, Gobbi C, Khullar V. The treatment of lower urinary tract symptoms in patients with multiple sclerosis: a systematic review. *Curr Urol Rep*. 2012 Oct;13(5):335-42. doi: 10.1007/s11934-012-0266-9.
15. Hoverd PA, Fowler CJ. Desmopressin In The Treatment Of Daytime Urinary Frequency In Patients With Multiple Sclerosis. *J Neurol Neurosurg Psychiatry*. 1998;65:778–80.
16. Kavia RB, de Ridder D, Costantinescu CS, et al. Randomized Controlled Trial Of Sativex To Treat Detrusor Overactivity In Multiple Sclerosis. *Mult Scler*. 2010;16(11):1349–59.
17. Capasso R, Aviello G, Borrelli F, et al. Inhibitory Effect Of Standardized Cannabis Sativa Extract And Its Ingredient Cannabidiol On Rat And Human Bladder Contractility. *Urology*. 2011;77 (4):1006e9–15.
18. Pryce G, Baker D. Potential Control Of Multiple Sclerosis By Cannabis And The Endocannabinoids System. *CNS Neurol Disord Drug Targets*. 2012,11.
19. Fijorbarck MV, Rijkhoff N, Petersen T, et al. Event Driven Electrical Stimulation Of The Dorsal Penile/Clitoral Nerve For Management Of Neurogenic Detrusor Overactivity In Multiple Sclerosis. *Neurourol Urodyn*. 2006;25:349–55.

20. Kabay SC, Yucel M, Kabay S. Acute Effect Of Posterior Tibial Nerve Stimulation On Neurogenic Detrusor Overactivity In Patients With Multiple Sclerosis: Urodynamic Study. *Urology*. 2008;71(4):641–5.
21. DeSeze M, Raibaut P, Gallien P, et al. Transcutaneous Posterior Tibial Nerve Stimulation For Treatment Of Overactive Bladder Syndrome In Multiple Sclerosis: A Multicenter Prospective Study. *Neurourol Urodyn*. 2011;30:306–11.
22. Fjorback MV, van Rey FM, Rijnkhoff MJ, et al. Electrical Stimulation Of Sacral Dermatomes In Multiple Sclerosis Patients With Neurogenic Detrusor Overactivity. *Neurourol Urodyn*. 2007;26:525–30.
23. Zchoval R, Pitha J, Medova E, et al. Augmentation cystoplasty in patients with multiple sclerosis. *Urol Int*. 2003;70:21–6.