

## Bibliografía

1. Ahmed RL, Thomas W, Yee D, Schmitz KH. Randomized Controlled Trial of Weight Training and Lymphedema in Breast Cancer Survivors. *J Clin Oncol*. 2006. 24:2765-2772.
2. Arias-Cuadrado A, Álvarez-Vázquez M J. Rehabilitación del linfedema; Actualización y protocolo. En: Servicio de Medicina Física y Rehabilitación del Hospital Comarcal de Valdeorras (Galicia) 2008.
3. Cavanaugh KM. Effects of Early Exercise on the Development of Lymphedema in Patients With Breast Cancer Treated With Axillary Lymph Node Dissection. *J Oncol Pract*. 2011 Mar; 7(2):89-93. doi: 10.1200/JOP.2010.000136.
4. Norman SA, Miller LT, Erikson HB, Norman MF, McCorkle R. Development and validation of a telephone questionnaire to characterize lymphedema in women treated for breast cancer. *Phys Ther* 2001. 81(6):1192-1205.
5. Panagiotou A, Nastoulis E, Bakirtzis D, Papadopoulos P, Tsonga Th, Fiska A et al. The role of exercise in the breast cancer related lymphedema. *Int J Med Med Sci*. 2011 Nov; 2(11):263-8.
6. Sagen A, Kåresen R, Risberg MA. Physical activity for the affected limb and arm lymphedema after breast cancer surgery. A prospective, randomized controlled trial with two years follow-up. *Acta Oncol*. 2009; 48(8):1102-10. doi: 10.3109/02841860903061683.
7. Schmitz KH, Ahmed RL, Troxel A, Cheville A, Smith R, Lewis-Grant L, et al. Weight lifting in women with breast-cancer-related lymphedema. *N Engl J Med*. 2009 Aug 13; 361(7):664-73.
8. Schmitz KH, Ahmed RL, Troxel AB, Cheville A, Lewis-Grant L, Smith R, et al. Weight lifting for women at risk for breast cancer-related lymphedema: a randomized trial. *JAMA*. 2010 Dec 22; 304(24):2699-705.
9. Ferreira DM, Goncalves IC, Rodrigues V, Goldner M. Fatores associados ao linfedema em pacientes com câncer de mama. *Rev. Bras. Ginecol. Obstet*. 2011; 33 (2):75-80.
10. Dr. Cernal Y, Pusic A, Mehrara BJ. Medidas preventivas para el linfedema: separando realidad de ficción. *J Am Coll Surg* 2011, 213(4): 543-551.
11. American Cancer Society, Inc. Consulta el 19 de enero de 2018. Disponible en: <http://www.cancer.org/treatment/treatmentsandsideeffects/physicalsideeffects/lymphedema/whateverywomanwithbreastcancershouldknow/lymphedema-with-breast-cancer-to-prevent-lymphedema>
12. Tewari N, Gill PG, Bochner MA, Kollias J. Comparison of volumen displacement versus circumferential arm measurements for lymphoedema: implications for the SNAC trial. *ANZ J. Surg*. 2008; 78: 889-893.
13. Hayes SC, Reul-Hirche H, Turner. Exercise and secondary lymphedema: safety, potential benefits, and research issues. *Med Sci Sports Exerc*. 2009 Mar; 41(3):483-9. doi: 10.1249/MSS.0b013e31818b98fb.
14. Johansson K, Lie E, Ekdahl CA. Randomized study comparing manual lymph drainage with secuencial pneumatic compression for treatment of postoperative arm lymphedema. *Lymphology*. 1998; 31: 56-64.