

Bibliografía

1. Gaby A. Multiple Sclerosis. *Glob Adv Health Med*. 2013 Jan;2(1):50-6. doi: 10.7453/gahmj.2013.2.1.009.
2. Horak F, Diener H. Cerebellar control of postural scaling and central set in stance. *J Neurophysiol*. 1994;72:479-493.
3. Nilsagard Y, Lundholm C, Denison E, et al. Predicting accidental falls in people with multiple sclerosis: a longitudinal study. *Clin Rehabil* 2009; 23: 259-269.
4. Mills RJ, Yap L, Young CA. Treatment for ataxia in multiple sclerosis. *Cochrane Database Syst Rev*. 2007 Jan 24;(1):CD005029.
5. McLoughlin J, Barr C1, Crotty M, Lord SR, Sturnieks DL. Association of Postural Sway with Disability Status and Cerebellar Dysfunction in People with Multiple Sclerosis: A Preliminary Study. *Int J MS Care*. 2015 May-Jun;17(3):146-51. doi: 10.7224/1537-2073.2014-003.
6. Quintern J, Immisch I, Albrecht H, Pöllmann W, Glasauer S, Straube A. Influence of visual and proprioceptive afferences on upper limb ataxia in patients with multiple sclerosis. *J Neurol Sci*. 1999 Feb 1;163(1):61-9.
7. Missaoui B, Thoumie P. How far do patients with sensory ataxia benefit from so-called "proprioceptive rehabilitation"? *Neurophysiol Clin*. 2009 Oct-Nov;39(4-5):229-33. doi: 10.1016/j.neucli.2009.07.002. Epub 2009 Aug 26.
8. Daley ML, Swank RL. Changes in postural control and vision induced by multiple sclerosis. *Agressologie*. 1983;24:327-329.
9. Prosperini L, Leonardi L, De Carli P, Mannocchi ML, Pozzilli C. Visuo-proprioceptive training reduces risk of falls in patients with multiple sclerosis. *Mult Scler*. 2010 Apr;16(4):491-9. doi: 10.1177/1352458509359923. Epub 2010 Feb 11.