

Bibliografía

1. Mudge S, Barber PA, Stott NS. Circuit-based rehabilitation improves gait endurance but not usual walking activity in chronic stroke: a randomized controlled trial. *Arch Phys Med Rehabil.* 2009 Dec;90(12):1989-96. doi: 10.1016/j.apmr.2009.07.015.
2. Dobkin BH. Strategies for stroke rehabilitation. *Lancet Neurol.* 2004;3:528-536.
3. de Bode S, Mathern GW, Bookheimer S, et al. Locomotor training remodels fMRI sensorimotor cortical activations in children after cerebral hemispherectomy. *Neurorehabil Neural Repair.* 2007;21:497-508.
4. Behrman AL, Harkema SJ. Locomotor training after human spinal cord injury: a series of case studies. *Phys Ther.* 2000;80:688-700.
5. Harkema SJ, Hurley SL, Patel UK, et al. Human lumbosacral spinal cord interprets loading during stepping. *J Neurophysiol.* 1997;77:797-811.
6. Beres-Jones JA, Harkema SJ. The human spinal cord interprets velocity-dependent afferent input during stepping. *Brain.* 2004;127:2232-2246.
7. Pang MY, Yang JF. The initiation of the swing phase in human infant stepping: importance of hip position and leg loading. *J Physiol (Lond).* 2000;528:389-404.
8. Moseley AM, Stark A, Cameron ID, et al. Treadmill training and body weight support for walking after stroke. *Cochrane Database Syst Rev.* 2005:CD002840.
9. Van Peppen RPS, Kwakkel G, Wood-Dauphinee S, et al. The impact of physical therapy on functional outcomes after stroke: what's the evidence? *Clin Rehabil.* 2004;18:833-862.
10. Polese JC, Ada L, Dean CM, Nascimento LR, Teixeira-Salmela LF. Treadmill training is effective for ambulatory adults with stroke: a systematic review. *J Physiother.* 2013 Jun;59(2):73-80. doi: 10.1016/S1836-9553(13)70159-0.
11. Charalambous CC, Bonilha HS, Kautz SA, Gregory CM, Bowden MG. Rehabilitating walking speed poststroke with treadmill-based interventions: a systematic review of randomized controlled trials. *Neurorehabil Neural Repair.* 2013 Oct;27(8):709-21. doi: 10.1177/1545968313491005. Epub 2013 Jun 13.
12. Lewek MD. The value of overground gait training for improving locomotion in individuals with chronic stroke. *J Neurol Phys Ther.* 2009 Dec;33(4):187-8. doi: 10.1097/NPT.0b013e3181c29aaa.
13. Ivey FM, Hafer-Macko CE, Macko RF. Exercise rehabilitation after stroke. *NeuroRx.* 2006;3:439-450.
14. Dickstein R. Rehabilitation of gait speed after stroke: a critical review of intervention approaches. *Neurorehabil Neural Repair.* 2008 Nov-Dec;22(6):649-60. doi: 10.1177/15459683080220060201.
15. Pang MY, Charlesworth SA, Lau RW, Chung RC. Using aerobic exercise to improve health outcomes and quality of life in stroke: evidence-based exercise prescription recommendations. *Cerebrovasc Dis.* 2013;35(1):7-22. doi: 10.1159/000346075. Epub 2013 Feb 14.
16. Pedersen BK, Saltin B. Exercise as medicine - evidence for prescribing exercise as therapy in 26 different chronic diseases. *Scand J Med Sci Sports.* 2015 Dec;25 Suppl 3:1-72. doi: 10.1111/sms.12581.