

Bibliografía

1. Bansil J, Bloch W, Gamper U, Kesselring J. Training in MS: influence of two different endurance training protocols (aquatic versus overland) on cytokine and neurotrophin concentrations during three week randomized controlled trial. *Mult Scler* (2013) 19:613–21. doi:10.1177/1352458512458605
2. Prosperini L, Piattella MC, Gianni C, Pantano P. Functional and structural brain plasticity enhanced by motor and cognitive rehabilitation in multiple sclerosis. *Neural Plast* (2015) 2015:481574. doi:10.1155/2015/481574.
3. Cramer SC, Sur M, Dobkin BH, O'Brien C, Sanger TD, Trojanowski JQ, et al. Harnessing neuroplasticity for clinical applications. *Brain* (2011) 134:1591–609. doi:10.1093/brain/awr039
4. Sterr A, Freivogel S. Motor-improvement following intensive training in low-functioning chronic hemiparesis. *Neurology*. 2003 Sep 23;61(6):842-4.
5. Lennon S, Ashburn A. The Bobath concept in stroke rehabilitation: a focus group study of the experienced physiotherapists' perspective. *Disabil Rehabil*. 2000 Oct 15;22(15):665-74.
6. Carmeli E, Peleg S, Bartur G, Elbo E, Vatine JJ. HandTutorTM Enhanced Hand Rehabilitation after Stroke — A Pilot Study. *Physiother. Res. Int.* 16 (2011) 191–200. DOI: 10.1002/pri.485.
7. Jezernik S, Colombo G, Keller T, Frueh H, Morari M. Robotic orthosis lokomat: a rehabilitation and research tool. *Neuromodulation*. 2003 Apr;6(2):108-15. doi: 10.1046/j.1525-1403.2003.03017.x.
8. Schwartz I, Sajin A, Moreh E, Fisher I, Neeb M, Forest A, Vaknin-Dembinsky A, Karusis D, Meiner Z. Robot-assisted gait training in multiple sclerosis patients: a randomized trial. *Mult Scler*. 2012 Jun;18(6):881-90. doi: 10.1177/1352458511431075.
9. Swinnen E, Beckwée D, Meeusen R, Baeyens JP, Kerckhofs E. Does robot-assisted gait rehabilitation improve balance in stroke patients? A systematic review. *Top Stroke Rehabil*. 2014 Mar-Apr;21(2):87-100. doi: 10.1310/tsr2102-87.
10. Straudi S, Manfredini F, Lamberti N, Zamboni P, Bernardi F, Marchetti G, Pinton P, Bonora M, Secchiero P, Tisato V, Volpato S, Basaglia N. The effectiveness of Robot-Assisted Gait Training versus conventional therapy on mobility in severely disabled progressive Multiple sclerosis patients (RAGTIME): study protocol for a randomized controlled trial. *Trials*. 2017 Feb 27;18(1):88. doi: 10.1186/s13063-017-1838-2.
11. Lefeber N, Swinnen E, Kerckhofs E. The immediate effects of robot-assistance on energy consumption and cardiorespiratory load during walking compared to walking without robot-assistance: a systematic review. *Disabil Rehabil Assist Technol*. 2017 Oct;12(7):657-671. doi: 10.1080/17483107.2016.1235620.
12. Mayr A, Quirbach E, Picelli A, Kofler M, Smania N, Saltuari L. Early robot-assisted gait retraining in non-ambulatory patients with stroke: a single blind randomized controlled trial. *Eur J Phys Rehabil Med*. 2018 Mar 29. doi: 10.23736/S1973-9087.18.04832-3.
13. Nam KY, Kim HJ, Kwon BS, Park JW, Lee HJ, Yoo A. Robot-assisted gait training (Lokomat) improves walking function and activity in people with spinal cord injury: a systematic review. *J Neuroeng Rehabil*. 2017 Mar 23;14(1):24. doi: 10.1186/s12984-017-0232-3.