

Bibliografía

1. Carr JH, Shepherd RB. Neurological rehabilitation. Optimizing motor performance. Oxford: Butterworth Heinemann; 1998.
2. Dean CM, Richards CL, Malouin F. Task-related circuit training improves performance of locomotor tasks in chronic stroke: a randomized, controlled pilot trial. *Arch Phys Med Rehabil.* 2000 Apr;81(4):409-17.
3. Tinson DJ. How stroke patients spend their days: an observational study of the treatment regime offered to patients in hospital with movement disorders following stroke. *Int Disabil Studies* 1989;11: 45-9.
4. Lerner-Frankiel MB, Vargas S, Brown MB, Krusell K, Schoneberger W. Functional community ambulation: what are your criteria? *Clin Manage* 1990;6:12-5.
5. Hill K, Ellis P, Bernhardt J, Maggs P, Hull S. Balance and mobility outcomes for stroke patients: a comprehensive audit. *Aust J Physiother* 1997;43:173-80.
6. Bustamante Valles K, Montes S, Madrigal Mde J, Burciaga A, Martínez ME, Johnson MJ. Technology-assisted stroke rehabilitation in Mexico: a pilot randomized trial comparing traditional therapy to circuit training in a Robot/technology-assisted therapy gym. *J Neuroeng Rehabil.* 2016 Sep 15;13(1):83. doi: 10.1186/s12984-016-0190-1.
7. English C, Hillier SL, Lynch EA. Circuit class therapy for improving mobility after stroke. *Cochrane Database Syst Rev.* 2017 Jun 2;6:CD007513. doi: 10.1002/14651858.CD007513.pub3.
8. Kaur G, English C, Hillier S. How physically active are people with stroke in physiotherapy sessions aimed at improving motor function? A systematic review. *Stroke Res Treatment.* 2012;2012:820673. doi: 10.115/2012/820673.
9. Intercollegiate Stroke Working Party. National clinical guidelines for stroke. 4E. London: Royal College of Physicians; 2012.
10. Bennett L, Luker J, English C, Hillier S. Stroke survivors' perspectives on two novel models of inpatient rehabilitation: seven-day a week individual therapy or five-day a week circuit class therapy. *Disabil Rehabil.* 2016 Jul;38(14):1397-406. doi: 10.3109/09638288.2015.1103788.