

Como conclusión, en función de los estudios publicados hasta la fecha, se recomienda pautar ejercicio aeróbico para los pacientes con distrofia facio-escápulo-humeral, si bien se necesitan más estudios para comprobar la eficacia del ejercicio asistido con estimulación eléctrica neuromuscular.

## Bibliografía

1. Voet NB, Bleijenberg G, Padberg GW, van Engelen BG, Geurts AC. Effect of aerobic exercise training and cognitive behavioural therapy on reduction of chronic fatigue in patients with facioscapulohumeral dystrophy: protocol of the FACTS-2-FSHD trial. *BMC Neurol*. 2010 Jun 30;10:56. doi: 10.1186/1471-2377-10-56.
2. Kissel JT: Facioscapulohumeral dystrophy. *Semin Neurol* 1999, 19:35-43.
3. Quarantelli M, Lanzillo R, Del VW, Mollica C, Prinster A, Iadicicco L, Iodice V, Santoro L: Modifications of brain tissue volumes in facioscapulohumeral dystrophy. *Neuroimage* 2006, 32:1237-1242.
4. van der Kooi EL, Kalkman JS, Lindeman E, Hendriks JC, van Engelen BG, Bleijenberg G, Padberg GW: Effects of training and albuterol on pain and fatigue in facioscapulohumeral muscular dystrophy. *J Neurol* 2007, 254:931-940.
5. Sackley C, Disler PB, Turner-Stokes L, Wade DT, Brittle N, Hoppitt T. Rehabilitation interventions for foot drop in neuromuscular disease. *Cochrane Database Syst Rev*. 2009 Jul 8;(3):CD003908. doi: 10.1002/14651858.CD003908.pub3.
6. Voet NB, van der Kooi EL, Riphagen II, Lindeman E, van Engelen BG, Geurts AC. Strength training and aerobic exercise training for muscle disease. *Cochrane Database Syst Rev*. 2013 Jul 9;(7):CD003907. doi: 10.1002/14651858.CD003907.pub4.
7. Brussee V, Tradif F, Tremblay JP. Muscle fibers of mdx mice are more vulnerable to exercise than those of normal mice. *Neuromuscul Disord* 1997;7:487-492.
8. Andersen G, Prahm KP, Dahlqvist JR, Citirak G, Vissing J. Aerobic training and postexercise protein in facioscapulohumeral muscular dystrophy: RCT study. *Neurology*. 2015 Aug 4;85(5):396-403. doi: 10.1212/WNL.0000000000001808.
9. Voet N, Bleijenberg G, Hendriks J, de Groot I, Padberg G, van Engelen B, Geurts A. Both aerobic exercise and cognitive-behavioral therapy reduce chronic fatigue in FSHD: an RCT. *Neurology*. 2014 Nov 18;83(21):1914-22. doi: 10.1212/WNL.0000000000001008.
10. Bankolé LC, Millet GY, Temesi J, Bachasson D, Ravelojaona M, Wuyam B, Verges S, Ponsot E, Antoine JC, Kadi F, Féasson L. Safety and efficacy of a 6-month home-based exercise program in patients with facioscapulohumeral muscular dystrophy: A randomized controlled trial. *Medicine (Baltimore)*. 2016 Aug;95(31):e4497. doi: 10.1097/MD.0000000000004497.
11. Colson SS, Benchortane M, Tanant V, Faghan JP, Fournier-Mehouas M, Benaïm C, Desnuelle C, Sacconi S. Neuromuscular electrical stimulation training: a safe and effective treatment for facioscapulohumeral muscular dystrophy patients. *Arch Phys Med Rehabil*. 2010 May;91(5):697-702. doi: 10.1016/j.apmr.2010.01.019.
12. Doix AM, Roeleveld K, Garcia J, Lahaut P, Tanant V, Fournier-Mehouas M, Desnuelle C, Colson SS, Sacconi S. Short-TERM Neuromuscular Electrical Stimulation Training of the Tibialis Anterior Did Not Improve Strength and Motor Function in Facioscapulohumeral Muscular Dystrophy Patients. *Am J Phys Med Rehabil*. 2017 Apr;96(4):e56-e63. doi: 10.1097/PHM.0000000000000705.