

Bibliografía

1. Currie CT. Urinary incontinence after stroke. *Br Med J (Clin Res Ed)* 1986 Nov;293(6558):1322-3. <http://www.ncbi.nlm.nih.gov/pubmed/3790967>
2. Codine PH, Pellissier J, Manderscheidt JC, Costa P, Enjalbert M, Perrigot M. Les troubles urinaires au cours des hémiplegies vasculaires. In: *Hémiplegie vasculaire et médecine de rééducation*. Pellissier J, ed. Paris, Masson, 1988, pp. 261-269.
3. Barer DH. Continence after stroke: useful predictor or goal of therapy? *Age Ageing* 1989 May;18(3):183-91. <http://www.ncbi.nlm.nih.gov/pubmed/2782216>
4. Nakayama H, Jørgensen HS, Pedersen PM, Raaschou HO, Olsen TS. Prevalence and risk factors of incontinence after stroke. The Copenhagen Stroke Study. 1997 Jan;28(1):58-62.
5. Khan Z, Hertenau J, Yang WC, Melman A, Leiter E. Predictive correlation of urodynamic dysfunction and brain injury after cerebrovascular accident. *J Urol* 1981 Jul;126(1):86-8.
6. Tsuchida S, Noto H, Yamaguchi O, Itoh M. Urodynamic studies on hemiplegic patients after cerebrovascular accident. *Urology* 1983 Mar;21(3):315-8.
7. Dumoulin C, Korner-Bitensky N, Tannenbaum C. Urinary incontinence after stroke: identification, assessment, and intervention by rehabilitation professionals in Canada. *Stroke*. 2007 Oct;38(10):2745-51.
8. Tibaek S, Dehlendorff C. Is Barthel Index a relevant measure for measuring prevalence of urinary incontinence in stroke patients? *Neurourol Urodyn*. 2012 Jan;31(1):44-9. doi: 10.1002/nau.21203.
9. Kohler M, Mayer H, Kesselring J, Saxer S. (Can) Not talk about it - Urinary incontinence from the point of view of stroke survivors: a qualitative study. *Scand J Caring Sci*. 2018 Mar;32(1):371-379. doi: 10.1111/scs.12471.
10. Tibaek S, Gard G, Klarskov P, Iversen HK, Dehlendorff C, Jensen R. Are activity limitations associated with lower urinary tract symptoms in stroke patients? A cross-sectional, clinical survey. *Scand J Urol Nephrol*. 2009;43(5):383-9. doi: 10.3109/00365590903013919.
11. Kohler M, Mayer H, Battocletti M, Kesselring J, Saxer S. Effectiveness of non-pharmacological interventions to promote urinary continence in stroke survivors – a systematic literature review. *Pflege*. 2016 Sep;29(5):235-245.
12. Booth J, Hagen S, McClurg D, et al. A feasibility study of transcutaneous posterior tibial nerve stimulation for bladder and bowel dysfunction in elderly adults in residential care. *J Am Med Dir Assoc*. 2013;14(4): 270–274.
13. Cooperberg MR, Stoller ML. Percutaneous neuromodulation. *Urol Clin North Am*. 2005;32(1):71–78, vii.
14. Guo ZF, Liu Y, Hu GH, Liu H, Xu YF. Transcutaneous electrical nerve stimulation in the treatment of patients with poststroke urinary incontinence. *Clin Interv Aging*. 2014 May 23;9:851-6. doi: 10.2147/CIA.S61084.
15. Liu Y, Xu G, Luo M, Teng HF. Effects of Transcutaneous Electrical Nerve Stimulation at Two Frequencies on Urinary Incontinence in Poststroke Patients: A Randomized Controlled Trial. *Am J Phys Med Rehabil*. 2016 Mar;95(3):183-93. doi: 10.1097/PHM.0000000000000360.
16. Shin DC, Shin SH, Lee MM, Lee KJ, Song CH. Pelvic floor muscle training for urinary incontinence in female stroke patients: a randomized, controlled and blinded trial. *Clin Rehabil*. 2016 Mar;30(3):259-67. doi: 10.1177/0269215515578695.
17. Resende AP, Stüpp L, Bernardes BT, Oliveira E, Castro RA, Girão MJ, Sartori MG 2012 Can hypopressive exercises provide additional benefits to pelvic floor muscle training in women with pelvic organ prolapse? *Neurourology and Urodynamics* 31: 121–125.
18. Mateus-Vasconcelos ECL, Ribeiro AM, Antônio FI, Brito LGO, Ferreira CHJ. Physiotherapy methods to facilitate pelvic floor muscle contraction: A systematic review. *Physiother Theory Pract*. 2018 Jun;34(6):420-432. doi: 10.1080/09593985.2017.1419520.
19. Mangera A, Chapple CR. Use of botulinum toxin in the treatment of lower urinary tract disorders. Current status. *Arch Esp Urol*. 2010 Dec;63(10):829-41.